

The effect of slurry composition on methane potential emissions from fattening pig slurries: a review of three nutrition assays

Salvador Calvet¹, Walter Antezana¹, Alba Cerisuelo², Fernando Estellés¹, Pablo Ferrer¹, Amanda Beccaccia³, Paloma García-Rebollar³, Carlos de Blas³

¹ Instituto de Ciencia y Tecnología Animal - Universitat Politècnica de València.

² Centro de Investigación y Tecnología Animal. Instituto Valenciano de Investigaciones Agrarias (IVIA).

³ Departamento de Producción Animal, E.T.S. Ingenieros Agrónomos, Universidad Politécnica de Madrid (UPM).

salcalsa@upvnet.upv.es

Palabras clave: methane emission, fattening pig, slurry composition, by products, nutritional strategy.

This study reviews the effects of pig slurry composition on the biochemical methane (CH₄) potential (B₀), using the information collected in three nutrition assays. A total of 84 animals were used to test the effect of 13 different diets. This included different types and sources of fibre (orange pulp and carob meal, study 1), different sources of protein (soybean meal, sunflower meal and wheat DDGS, study 2) and different inclusion levels of fat and fibre (calcium soap of palm fatty acids distillate and orange pulp, respectively, study 3). Faeces, urine and slurry production and composition were determined, as well as B₀ from slurry in an in-vitro assay during 100 days. Potential emissions per animal (CH₄-animal) were also calculated according to the daily excretion of volatile solids (VS). Correlations and multiple linear regression analysis were obtained to explain the factors affecting the emissions. B₀ ranged between 256 and 430 g CH₄ per gram of VS and was conditioned by the type of diet. Slurry excretion was affected by diet and originated major variations of potential CH₄ emitted per animal and day (66-144 L CH₄/day). Faeces and slurry pH were positively and strongly correlated with B₀ (r=0.72 and r=0.62, respectively, p<0.05). Nitrogen content of faeces was positively correlated with B₀ (0.61, p<0.05) and CH₄ produced per animal (r=0.77, p<0.01). No other relevant correlations with effluent characteristics were found. The following significant regression models were obtained: B₀ (mL/g VS)= -458+119.4*pH (R²=0.52) and CH₄-animal (L/day) = 78.5 - 0.82 * Urine dry matter (g/kg) + 0.39 crude protein in faeces (g/kg) (R² = 0.75). Although no clear relationship was found between effluent composition and potential CH₄ emissions, diet composition was a critical factor. The inclusion of different types of fibre and fat changed potential emission per animal in a two-fold range because of changes in B₀ of slurry and the amount of excreta produced.